

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/11, C07H 21/00, A61K 31/70,</b> <b>C07K 14/025, C12N 9/00</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/13071</b>
			<b>(43) International Publication Date:</b> 18 March 1999 (18.03.99)
<b>(21) International Application Number:</b> PCT/US98/18320			<b>(74) Agent:</b> ALTMAN, Daniel, E.; Knobbe, Martens, Olson & Bear, 16th floor, 620 Newport Center Drive, Newport Beach, CA 92660 (US).
<b>(22) International Filing Date:</b> 3 September 1998 (03.09.98)			
<b>(30) Priority Data:</b> 08/929,140 5 September 1997 (05.09.97) US			<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
<b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application</b> US 08/929,140 (CIP) Filed on 5 September 1997 (05.09.97)			<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(71) Applicant (for all designated States except US):</b> THE GOV- ERNMENT OF THE UNITED STATES OF AMERICA, as represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; Suite 325, 6011 Executive Boulevard, Rockville, MD 20852 (US).			
<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> DIPAOLO, Joseph [US/US]; 6605 Melody Lane, Bethesda, MD 20817 (US). ALVAREZ-SALAS, Luis [MX/US]; 912 Old Georgetown Road, Bethesda, MD 20814 (US).			
<b>(54) Title:</b> HUMAN PAPILLOMA VIRUS INHIBITION BY ANTI-SENSE OLIGONUCLEOTIDES			
<b>(57) Abstract</b> <p>Antisense oligonucleotides and analogs thereof having phosphorothioate backbone structure and sequences complementary to nucleotides contained with residues 415 to 445 of human papilloma virus 16 (HPV-16). These antisense oligonucleotides and oligonucleotide analogs are used to inhibit HPV-16 expression and to inhibit the growth of cervical tumors.</p>			
<p>The diagram illustrates the structure of the HPV-16 E6 and E7 genes and the binding sites for antisense oligonucleotides. The top part shows a linear representation of the E6 and E7 genes, with E6 starting at position 104 (ATG) and E7 starting at position 562 (ATG). Below this, a horizontal line represents the E6 and E7 genes, with a 'cop' (copy) label at position 97 and a vertical line at position 867. The middle part shows a detailed view of the E6(II)E7 (Minor) gene, with a vertical line at position 226 and a vertical line at position 526. The bottom part shows a detailed view of the E6(I)E7 (Major) gene, with a vertical line at position 226 and a vertical line at position 409. The bottom part also shows the sequence UAACUGUCAAAGCCACUGUGUCCUGAAGAA, which is complementary to the sequence 415 to 445 of the HPV-16 E6 and E7 genes.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## HUMAN PAPILLOMA VIRUS INHIBITION BY ANTI-SENSE OLIGONUCLEOTIDES

### Field of the Invention

The present invention relates to the use of antisense oligonucleotides to inhibit a Human Papilloma virus (HPV), and specifically relates to use of antisense oligonucleotides specific for nucleotides 415 to 445 of the DNA sequence of HPV-16.

### Background of the Invention

Papilloma viruses are small DNA viruses that induce the hyperproliferation of epithelial cells. Approximately 70 different genotypes of human papilloma virus (HPV) have been isolated. Some HPV genotypes (e.g., 1, 2, 4, and 7) have been associated with human benign squamous papillomas (warts and condylomas) and others (e.g., 16 and 18) have been associated with human neoplastic and preneoplastic lesions (DiPaolo, et al., 1993, *Crit. Rev. Oncogen.* 4:337-360).

HPV-16 has been associated with a variety of clinical conditions in both women and men. In women, HPV-16 is frequently associated with latent infections, benign and premalignant cervical lesions (dysplasias/CIN) and half of invasive cervical carcinomas. Cervical cancer, which kills at least 500,000 women worldwide each year, proceeds through progressive cellular changes from benign condylomata to high-grade dysplasias/CIN before developing into an invasive cancer. In men, HPV-16 is associated with subclinical macular or clinical papular lesions. One such lesion, Penile Bowenoid papulosis, resembles cervical carcinoma *in situ*. Detection and treatment of these lesions costs over five billion health care dollars annually in the United States.

HPV-16 has been associated with over half of the invasive cervical carcinomas diagnosed worldwide and with many cell lines derived from cervical carcinomas. HPV-16 expression causes benign proliferation and efficiently immortalizes cultured human epithelial cells, including cervical keratinocytes (DiPaolo, et al., 1993, *Crit. Rev. Oncogen.* 4:337-360; Zur Hausen & de Villiers, 1994, *Annu. Rev. Microbiol.* 48:427-447; Schiffman, 1995, *J. Natl. Cancer Inst.* 87:1345-1347). Two HPV-16 genes, E6 and E7, and their gene products are required to immortalize human keratinocytes and are a hallmark of cervical carcinoma (Hawley-Nelson et al., 1989, *EMBO J.* 8:3905-3910; Phelps et al., 1988, *Cell* 53:539-547; Viallet et al., 1994, *Exp. Cell Res.* 212:36-41; Yokoyama et al., *Obstet. Gynecol.* 83:197-204). The E6 and E7 proteins bind to other gene products (p53 and Rb tumor suppressors) to disrupt control of cell division and proliferation, leading to transformation (Scheffner et al., 1990, *Cell* 63:1129-1136; Zerbass et al., *J. Virol.* 69:6389-6399)

Surgery is commonly used for treatment of high-grade lesions due to the lack of effective alternatives. Cervical laser ablation therapy, however, does not in the long term influence the natural history of cervical human papillomavirus-associated diseases in women. Interferons have not proved an effective antiviral or anticancer treatment. Chemotherapy (e.g., cisplatin, alone or combined with other chemotherapy agents such as 5-fluorouracil) has generally not proved to be effective in treatment of many cervical cancers. Moreover, most chemotherapeutic agents are cytotoxic, leading to toxic side effects and the development of multiple drug resistance. Therefore, there

is a need for reagents than can specifically inhibit the growth of HPV-associated tumor cells, while avoiding serious toxic reactions.

HPV-specific treatments in the form of cleavage of HPV-specific RNA with ribozymes and inhibition by HPV-specific antisense oligonucleotides have been suggested (PCT International Patent Application WO 95/31552; DiPaolo, et al., 1993, *Crit. Rev. Oncogen.* 4:337-360; Steele, et al., 1993, *Cancer Res.* 53:2330; Storey, et al., 1991, *Nuc. Acids Res.* 19(15):4109). Ribozymes are small catalytic RNA molecules that can hybridize to and cleave a complementary RNA target (Cech, 1988, *JAMA* 260:3030-3034). Ribozymes having a "hairpin" motif have been found to be more efficient than the "hammerhead" motif (Hampel & Tritz, 1989, *Biochem.* 28:4929-4933; Hampel, et al., 1990, *Nuc. Acids Res.* 18:299-304) and "hairpin" ribozymes have been used to cleave viral targets, including the human immunodeficiency virus (HIV-1) and HPV (Ojwang, et al., 1992, *Proc. Natl. Acad. Sci. USA* 89, 10802-10806; Yu, et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:6340-6344; PCT International Patent Application WO 95/31552).

Antisense RNA and oligonucleotides hybridize to complementary mRNA, thus blocking translation and promoting the activity of endogenous RNase H to cleave the mRNA (Walder, 1988, *Genes Dev.* 2:502-504; Cohen, 1991, *Antisense Res. Dev.* 1:191-193). Although antisense RNA and oligonucleotides should be specific for their target sequence, nonspecific toxicity has been observed (Henry et al., 1997, *Toxicol.* 116:77-88; Henry et al., 1997, *Anticancer Drug Des.* 12:1-14). First-generation antisense phosphorothioates, whose nucleotide backbones carry sulfur atoms to slow intracellular degradation were often ineffective because of their inability to enter cells or to complement the target mRNA, but improved second generation phosphorothioate antisense therapies, referred to as "mixed backbone oligonucleotides" and "end-modified chimerics" that carry 2'-O-methylribonucleoside moieties have proven effective in clinical trials (Monia et al., 1996, *Nature Medicine* 6:668-675; Roush, 1997, *Science* 276:1192-1193; Agrawal et al., 1997, *Proc. Natl. Acad. Sci. USA* 94:2620-2625; Agrawal, 1996, *TIBTECH* 14:3-14).

Antisense inhibition of HPV-18 E6 and E7 expression in cell lines (C4-1 and HeLa) resulted in a significant decrease in growth rate with continuous addition of oligonucleotide (Steele, et al., 1993, *Cancer Res.* 53:2330-2337). Similar results have been observed in cells transfected with recombinant vectors (von Knebel Doeberitz & Gissmann, 1987, *Hamatol. Bluttransfus.* 31:377-279; Hamada et al., 1996, *Gynecol. Oncol.* 63:219-227).

The present invention discloses oligonucleotide sequences and methods of antisense therapy using antisense oligonucleotides defined by selected HPV-16 complementary sequences.

#### Summary of the Invention

According to the present invention, antisense oligonucleotides that specifically bind to a human papilloma virus-16 (HPV-16) sequence include sequences complementary to viral sequences between viral nucleotide 415 and 445.

One aspect of the present invention relates to analogs of antisense oligonucleotides comprising oligonucleotide sequences complementary to SEQ ID NO: 2 or SEQ ID NO: 3, wherein the analogs are phosphorothioate antisense oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond, mixed backbone antisense oligonucleotides in which at least one phosphodiester bond is

replaced with a phosphorothioate bond and at least one phosphodiester bond is replaced with a 2'-O-methylnucleoside phosphodiester bond, end-modified analogs in which at least one end has a 2'-O-methylnucleotide moiety, methylphosphonates, phosphoramidites, phosphorodithioates, or N3'→P5'-phosphoramidites.

Another aspect of the present invention relates to analogs of antisense oligonucleotides comprising  
5 sequences of SEQ ID NO:6, SEQ ID NO:7, or SEQ ID NO:17, wherein the analogs are phosphorothioate oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond, mixed backbone oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond and at least one phosphodiester bond is replaced with a 2'-O-methylnucleoside phosphodiester bond, end-modified oligonucleotides in which at least one end has a 2'-O-methylnucleotide moiety, methylphosphonates, phosphoramidites,  
10 phosphorodithioates, or N3'→P5'-phosphoramidites.

Another aspect relates to analogs of antisense oligoribonucleotides of SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:16, wherein the analogs are oligoribonucleotide phosphorothioates, 2'-O-alkyl oligoribonucleotide phosphorothioates or 2'-O-methylribonucleotide methylphosphonates. Yet another aspect of the invention relates to an antisense therapeutic composition comprising an oligonucleotide sequence of SEQ ID NO: 9 and a pharmaceutically  
15 acceptable carrier.

Therapeutic compositions include any of the foregoing analogs and a pharmaceutically acceptable carrier. The therapeutic compositions can also include a ribozyme containing sequences complementary to SEQ ID NO: 2 or SEQ ID NO: 3.

Yet one more aspect of the present invention relates to a method of preventing transformation of a living  
20 cell by HPV. The method includes providing an antisense therapeutic composition as is described above, providing a living cell capable of being transformed by HPV, transfecting the living cell with the antisense therapeutic composition, and maintaining the viability of the living cell for sufficient time to inhibit expression of HPV gene E6.

Still another aspect of the present invention is a method of preventing transformation of a living cell by HPV. This method includes providing one or more antisense oligonucleotides having sequences complementary to SEQ  
25 ID NO: 2 or SEQ ID NO: 3, providing a living cell capable of being transformed by HPV, transfecting the one or more antisense oligonucleotide into the living cell, and maintaining the viability of the living cell for sufficient time to inhibit expression of HPV gene E6. The living cell can be a human keratinocyte, a human cervical cell, or other living cell.

An additional aspect of the invention relates to a method for inhibiting expression of HPV gene E6 in a living cell comprising the steps of: providing one or more antisense oligonucleotides having sequences complementary to  
30 SEQ ID NO: 2 or SEQ ID NO: 3; providing a biological sample comprising living cells capable of being infected with HPV; transfecting said one or more antisense oligonucleotides into said living cells; and maintaining the viability of said living cells for sufficient time to allow inhibition of HPV E6 gene expression to occur. The method can also include repeating the transfecting and maintaining steps. The living cell can be a human keratinocyte, a human cervical cell, or other living cell. The step of providing antisense oligonucleotides can include administering antisense  
35 oligonucleotides to a living organism by subcutaneous (s.c.), intraperitoneal (i.p.) or intravenous (i.v.) injection, or by painting the antisense oligonucleotides onto the biological sample *in situ*.

An additional aspect of the invention relates to a method of inhibiting expression of HPV gene E6 in a living cell. This method includes the steps of: providing one or more antisense oligonucleotide analogs having sequences of SEQ ID NO:6, SEQ ID NO:7, or SEQ ID NO:17, wherein the one or more antisense oligonucleotide analogs are phosphorothioate oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond, mixed backbone oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond and at least one phosphodiester bond is replaced with a 2'-O-methylnucleoside phosphodiester bond, in which at least one end has a 2'-O-methylnucleotide moiety, methylphosphonates, phosphoramidites, phosphorodithioates, or N3'→P5'-phosphoramidites; providing a biological sample comprising living cells capable of being infected with HPV; transfecting said antisense oligonucleotide analogs into said living cells; and maintaining the viability of said living cells for sufficient time to allow inhibition of HPV E6 gene expression to occur.

The present invention also includes another aspect which is a method for inhibiting expression of HPV gene E6 in a living cell comprising the steps of: providing one or more antisense oligoribonucleotide analogs of oligoribonucleotides of SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:16, wherein the analogs are oligoribonucleotide phosphorothioates, 2'-O-alkyl oligoribonucleotide phosphorothioates or 2'-O-methylribonucleotide methylphosphonates; providing a biological sample comprising living cells capable of being infected with HPV; transfecting said antisense oligoribonucleotide analogs into said living cells; and maintaining the viability of said living cells for sufficient time to allow inhibition of HPV E6 gene expression to occur.

The present invention also provides a method for inhibiting the growth of cervical tumors, comprising the step of contacting the tumors with one or more analogs of antisense oligonucleotides comprising oligonucleotide sequences complementary to SEQ ID NO:2 or SEQ ID NO:3, wherein the one or more analogs are phosphorothioate antisense oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond, mixed backbone antisense oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond and at least one phosphodiester bond is replaced with a 2'-O-methylnucleoside phosphodiester bond, end-modified analogs in which at least one end has a 2'-O-methylnucleotide moiety, methylphosphonates, phosphoramidites, phosphorodithioates, or N3'→P5'-phosphoramidites.

Yet another embodiment of the invention is the use of one or more analogs of antisense oligonucleotides comprising oligonucleotide sequences complementary to SEQ ID NO:2 or SEQ ID NO:3, wherein the one or more analogs are phosphorothioate antisense oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond, mixed backbone antisense oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond and at least one phosphodiester bond is replaced with a 2'-O-methylnucleoside phosphodiester bond, end-modified analogs in which at least one end has a 2'-O-methylnucleotide moiety, methylphosphonates, phosphoramidites, phosphorodithioates, or N3'→P5'-phosphoramidites, for treatment of a cervical tumor.

#### Brief Description of the Drawings

FIG. 1 is a diagram of the HPV-16 E6/E7 target sites showing the overlaps in the E6/E7 mRNA and the ATG sites of E6 (at nt 104) and of E7 (at nt 562), the primary mRNA transcript having a cap at nt 97 and terminating

at nt 867, a minor form of processed E6/E7 mRNA ("E6(II)E7") spliced at nt 226 and 526, and a major form of processed E6/E7 mRNA ("E6(I)E7") spliced at nt 226 and 409, and the relative position of the antisense target sequence between nt 415 and 445. The complete HPV-16 sequence (GenBank Accession No. K02718) is presented in SEQ ID NO:1 and the antisense target sequence between nt 415-445 occurs therein.

5           FIG. 2 is a diagram of a hairpin ribozyme for cleavage of HPV-16 E6/E7 mRNA with an optimized helix 1, comprising 8 bp, designed to cleave HPV-16 after position 434 ("cleavage site" indicated by the diagonal line), showing the sequences of the ribozyme (in capital letters) and the substrate (in lower case letters), the regions of base pairing between target substrate (nt 430-445) and ribozyme ("helix 1" and "helix 2"), and the regions of base pairing predicted for the hairpin portion of the ribozyme ("helix 3" and "helix 4").

10           FIG. 3 shows a gel of ribozyme products of the R434 and R434i ribozymes produced by in vitro transcription and incubated with substrate RNA at a 1:2 molar ratio; the arrows at the right show the positions of the uncleaved substrate, and the 3' and 5' fragments produced by substrate cleavage.

            FIG. 4 diagrams the cytomegalovirus promoter/enhancer ("CMV") expression plasmids containing the HPV-16 E6/E7 genes only ("pCR16HH"), and with the *cis*-acting R434 ribozyme or R434i inactive ribozyme coding sequences ("pCR16E6/E7Rz" and "pCR16E6/E7Rzi" respectively).

15           FIG. 5 shows a graph of cell growth of normal human keratinocytes (HKc) (cell count on the Y-axis) over 2 to 7 days in culture (X-axis) for HKc transfected with the control plasmid, pCR16HH (—○—), the active ribozyme construct, pCR16E6/E7RZ (—□—) and the inactive ribozyme construct, pCR16E6/E7RZi (—△—).

            FIG. 6 shows an agarose gel separation of the products of a RT-PCR assay specific for HPV-16 E6/E7 mRNA. HKc were transfected with the E6/E7 construct without any ribozyme sequences (pCR16HH), the active ribozyme construct (pCR16E6/E7RZ), and the inactive ribozyme construct (pCR16E6/E7RZi); the negative "Control" is a RT-PCR reaction run without reverse transcriptase. The lower panel shows that all cells produced the 661 bp  $\beta$ -actin band; the upper panel shows the 492 bp uncleaved E6/E7 transcript band and an internal control band of 326 bp.

25           FIG. 7 is a graph showing the cell counts (Y-axis) for HKc transfected with the E6/E7 construct without any ribozyme sequences (pCR16HH, left bar), the active ribozyme construct (pCR16E6/E7RZ, middle bar), and the inactive ribozyme construct (pCR16E6/E7RZi, right bar) at 8 weeks of growth following transfection.

            FIG. 8 is a graph showing inhibition of growth of CasKi cervical tumor cells in culture. Anti-E6, M4 and M7 oligodeoxynucleotides were phosphorothioated and 50  $\mu$ M was applied to CasKi cervical carcinoma cells for 72 hours. Cell growth was estimated by the colorimetric quantitation of Br-dUTP incorporation. Similar results were obtained for QGU cervical carcinoma cells.

#### Detailed Description of the Preferred Embodiments

            The present invention provides antisense oligonucleotides, oligoribonucleotides and analogs thereof, for inhibiting the expression of HPV-16 E6 and E7 genes which are necessary for viral replication. Because the E6 and E7 genes overlap and produce mRNA molecules that overlap the two genes (as shown in FIG. 1), the E6 and E7 genes together and their mRNA overlapping transcripts will be generally referred to herein as E6/E7 genes and E6/E7

mRNA, respectively. These antisense molecules bind to E6/E7 mRNA in the cell, prevent mRNA translation and promote mRNA degradation by intracellular RNase H. These molecules inhibit the growth of cervical tumor cell lines *in vitro*, and inhibit cervical tumor growth *in vivo* in nude mice.

In the course of characterizing ribozymes that cleave E6/E7 mRNA, ribozymes that are inactive because of changes to the hairpin structure of the ribozyme were found to inhibit cell proliferation *in vitro* showing that the ribozymes were, in fact, acting at least in part as antisense inhibitors. That is, even in the absence of ribozyme-mediated cleavage of E6/E7 mRNA, the introduced ribozyme sequences that contain antisense sequences directed to E6/E7 target sequences were capable of limiting the amount of full-length E6/E7 transcripts. These antisense sequences are the basis for antisense oligonucleotides having modified backbone structure for use as antisense therapeutics.

#### Ribozyme Constructs and Activity

Potential hairpin ribozyme target sites were identified in the HPV-16 gene sequence for E6/E7 contained within the complete HPV-16 sequence (SEQ ID NO:1), by using a computer program (e.g., GCG DNA Analysis Package, Genetics Computer Group, Madison, WI) to search for 5'-GUC-3' motifs. Several potential cleavage sites were identified in E6 (nt 419, 434, 491, 503 and 514) and E7 (nt 679), and synthetic ribozymes covalently linked to sequences complementary to 15 or 16 nt sequences surrounding the identified potential cleavage sites were produced. The initially synthesized ribozymes were based on the structures of the negative strand of satellite RNA from the tobacco ringspot virus (-sTRSV) (Hampel & Tritz, 1989, *Biochem.* 28:4929-4933; Haseloff & Gerlach, 1989, *Gene* 82:43-52). Of these synthetic ribozymes, only those having sequences complementary to the E6 sites at nt 419 and 434 were found to significantly cleave substrate RNA (greater than 60% when mixed at a 1:2 molar ratio of ribozyme to substrate RNA). Improved ribozymes were then synthesized for these particular target sequences, in which modifications were introduced in helix 4 (based on the findings of Anderson et al., 1994, *Nucl. Acids Res.* 22:1096-1100). An improved ribozyme structure having a sequence complementary to target sequence of nt 430-445 of E6 mRNA is shown in FIG. 2. Using cell free reactions to optimize the length of helix 1 (in the hybrid formed between the target sequence and the ribozyme complementary sequence), the optimum sequences in the target were those corresponding to the HPV-16 sequence from nt 415-429 (UAACUGUCAAAGCC; SEQ ID NO:2) and nt 430-445 (ACUGUGUCCUGAAGAA; SEQ ID NO:3). Based on these target sequences, antisense oligonucleotides interacting with these target sequences have the following sequences: GGCUUUAGAAGUUA (SEQ ID NO:4) and UUCUUCAGAGAACAGU (SEQ ID NO:5), for antisense RNA complementary to nt 415-429 and to nt 430-445 of E6, respectively; and GGCTTTTAGAAGTTA (SEQ ID NO:6) and TTCTTCAGAGAACAGT (SEQ ID NO:7), for antisense DNA complementary to nt 415-429 and to nt 430-445 of E6, respectively.

The ribozyme coding sequences were synthesized and cloned using standard procedures. The coding sequences were synthesized in an automated DNA synthesizer (Expedite 8900, Perseptive Biosystems, Framingham, MA) and cloned into a plasmid (pBluescript KS vector, Strategene, La Jolla, CA). Ribozyme coding sequences were cloned *in cis* to complete E6/E7 gene sequences in another plasmid; the E6/E7 and ribozyme sequences were PCR amplified using standard procedures and cloned into another vector (pCR3.1, Invitrogen Corp., San Diego, CA) to



produce plasmids capable of transcribing ribozymes and target sites in *cis*. Plasmids were linearized and purified using standard procedures (restriction digestion and QIAquick column, Qiagene Inc., Chatsworth, CA) and 1  $\mu$ g of linear DNA template was incubated with T3 or T7 RNA polymerase, rNTP and  $\alpha$ -<sup>32</sup>P-UTP (Amersham Life Sciences, Arlington Heights, IL) to produce ribozyme and/or RNA substrate using standard procedures (e.g., as provided by  
 5 Ambion Inc., Austin, TX). Target RNA was gel purified (6% polyacrylamide/7M urea gel) before use by standard methods.

The active ribozyme that is specific for nt 430-445, designated R434, consists of the ribozyme sequence  
 d i a g r a m m e d i n F I G . 2 ( S E Q I D N O : 8 ;  
 UUCUUCAGAGAACAGUACCAGAGAAACACACGGACUUCGGUCCGUGGUUAUUAUACCUGGUA). An inactive ribozyme in  
 10 which the A<sub>24</sub>, A<sub>25</sub> and A<sub>28</sub> residues of SEQ ID NO:8 have been replaced with C, G and U, respectively, is referred to as R434i and consists of the ribozyme sequence of SEQ ID NO:9  
 (UUCUUCAGAGAACAGUACCAGAGCGUCACACGGACUUCGGUCCGUGGUUAUUAUACCUGGUA). The active ribozyme that  
 contains sequences complementary to E6 nt 415-430 is referred to as R419 and consists of  
 GGCUUUUAGAAGUUAACCAGAGAAACACACGGACUUCGGUCCGUGGUUAUUAUACCUGGUA (SEQ ID NO:10).

#### In Vitro Ribozyme Activity

Ribozyme activity was measured *in vitro* initially at 37°C in reaction buffer (40 mM Tris-HCl, pH 7.5, 12 mM MgCl<sub>2</sub>, 2 mM spermidine) containing 25 nM <sup>32</sup>P-labeled ribozyme and 50 nM <sup>32</sup>P-labeled substrate for 60 min; complete characterization was done using similar reactions except that 1 nM <sup>32</sup>P-labeled ribozyme and 30 nM <sup>32</sup>P-labeled substrate (1:30 molar ratio) were incubated for 180 min. Ribozyme expression from linear or covalently-  
 20 closed templates was accomplished by incubating 1  $\mu$ l of an *in vitro* transcription reaction (described above) with 10<sup>6</sup> cpm of <sup>32</sup>P-labeled target RNA in 10  $\mu$ l of reaction mixture. Reactions were stopped by freezing on dry ice; samples were denatured in loading buffer (80% formamide, 0.01% bromophenol blue, 0.01% xylene cyanol) at 65°C for 10 min and separated by gel electrophoresis (6% polyacrylamide/7M urea gel) using standard methods. Dried gels were exposed to radiographic film (BM-2, Kodak Corp., Rochester, NY) and bands intensities indicative of  
 25 uncleaved and/or cleaved RNA substrate were quantified using a Phosphorimager 425 (Molecular Dynamics, Sunnyvale, CA).

Ribozymes R419 and R434 were active in the *in vitro* ribozyme reactions in which there was 30-fold excess substrate. The R419 ribozyme had a calculated K<sub>m</sub> of 0.098  $\mu$ M and a k<sub>CAT</sub> of 0.18 min<sup>-1</sup>; the R434 ribozyme had a calculated K<sub>m</sub> of 0.021  $\mu$ M and a k<sub>CAT</sub> of 0.08 min<sup>-1</sup>. The catalytic efficiency (k<sub>CAT</sub>/K<sub>m</sub>) of R434 (3.81  $\mu$ M<sup>-1</sup>) was  
 30 twice as high as that of R419 (1.84  $\mu$ M<sup>-1</sup>). These results show that the antisense oligonucleotide sequences (SEQ ID NO:4 and SEQ ID NO:5) contained in ribozymes R419 and R434 are capable of specifically binding to the HPV-16 target sequences. By analogy, DNA antisense oligonucleotide sequences (SEQ ID NO:6 and SEQ ID NO:7) are equally capable of binding to the HPV-16 target sequences in E6.

The R434i ribozyme is changed in the ribozyme hairpin but contains the same target-recognition site as the  
 35 R434 ribozyme. These changes abolished catalytic activity of the R434i ribozyme *in vitro* as shown in FIG. 3, in

which aliquots of an *in vitro* reaction were analyzed at 0, 15, 30 and 60 min. for R434 and R434i ribozyme reactions.

#### In Vivo Ribozyme Activity in Cultured Human Keratinocytes

The R434 and R434i ribozyme sequences were cloned into plasmids for transfection into cells for measurement of *in vivo* activity. Plasmid pCR16-E6/E7RZ contains the E6/E7 gene sequences linked in *cis* to the 5' end of enzymatically-active ribozyme R434 sequence, and plasmid pCR16-E6/E7RZi contains the E6/E7 gene sequences linked in *cis* to the 5' end of enzymatically-inactive ribozyme R434i sequence. In both plasmids, the constructs were under the control of the cytomegalovirus (CMV) promoter/enhancer sequences as diagramed in FIG. 4. Control plasmids ("pCR16HH") contained the E6/E7 gene sequences under the control of the CMV promoter/enhancer sequences but without any ribozyme sequences.

Normal human keratinocytes (HKc) from neonatal foreskins were cultured in MCDB151-LB medium using standard methods (Pirisi et al., 1988, *Carcinogen*. 9:1573-1579) and transfected with 10  $\mu$ g of plasmid DNA using standard lipofection methods (Lipofectin, Life Technologies Inc., Gaithersburg, MD; Alvarez-Salas et al., 1995, *Cancer Lett.* 91:85-92). Transfected cells were grown in the presence of 200  $\mu$ g/ml of G418 for two weeks (or four days for immortalization studies) and growth rates were determined in standard six-well plates ( $10^6$  cells/well) in triplicate; cells were counted at the end of the incubation period (Coulter Counter ZM, Coulter Electronics Inc., Hialeah, FL).

Following transfection of HKc with the plasmid constructs, cell growth was assayed at 2 to 7 days post-transfection (shown in FIG. 5). HKc transfected with the active ribozyme construct (pCR16E6/E7RZ,  $-\square-$ ) grew significantly slower than cells transfected with the control plasmid (pCR16HH,  $-\bigcirc-$ ) or the inactive ribozyme (pCR16E6/E7RZi,  $-\triangle-$ ) construct. The latter two were capable of expressing the E6/E7 gene products, whereas the active ribozyme would have limited E6/E7 gene expression. To confirm this, a reverse transcription-polymerase chain reaction (RT-PCR) assay was performed to detect the products of ribozyme cleavage.

The RT-PCR assay was performed as follows. Total RNA was purified from the cultured cells (Rneasy Kit, Qiagen) using standard methods. HPV-16 E6/E7 cDNA was produced from 1  $\mu$ g of total RNA using standard methods (Superscript II One Shot Kit, Life Technologies). To produce differential sized bands for cleaved and uncleaved E6/E7 mRNA, the upper PCR primers were SEQ ID NO:11 (CAGCAATACAACAAACCG) and SEQ ID NO:12 (CACGTAGAAACCCAGC), flanking the R434 target site (nt 371-388 and nt 537-554, respectively), and the lower primer was SEQ ID NO:13 (TAGATTATGGTTTCTGAGAACA), hybridizing in the E7 gene (nt 862-841). Standard PCR conditions were used (as supplied by Strategene, La Jolla, CA) with the following times and temperatures. The first strand cDNA was synthesized for 30 min at 45°C, followed by denaturation (92°C for 2 min) and 35 PCR cycles of: denaturation (92°C for 1 min), hybridization of primers (45°C for 45 sec) and polymerization (72°C for 1 min). This PCR amplification produced two products: an uncleaved product of 492 bp (amplified by SEQ ID NO: 11 and SEQ ID NO:13) and an internal control product of 326 bp (amplified by SEQ ID NO: 12 and SEQ ID NO:13). A control PCR reaction under the same conditions but with primers specific for an endogenous  $\beta$ -actin gene (SEQ ID NO:14: TGACGGGGTCAACCACACTGTGCCCCATCTA, and SEQ ID NO:15:

CTAAGAAGCATTTCGCGGTGGACGATGGAGGG) was used as a control to produce a band of 661 bp. Amplified products were separated on a 1.5% agarose gel and visualized with long-wave UV after ethidium bromide staining.

As shown in FIG. 6, all of the transfected cells produced the 661 bp  $\beta$ -actin control band. The control transfected (pCR16HH) cells and the inactive ribozyme construct (pCR16E6/E7RZi) transfected cells produced both the 492 bp and 326 bp products showing the presence of full-length E6/E7 transcripts, whereas the active ribozyme construct (pCR16E6/E7RZ) produced no detectable amount of the 492 bp band, indicating cleavage. These results support the finding that the decreased growth rate of the HKc transfected with the active ribozyme construct (pCR16E6/E7RZ) was due to the inhibition of E6/E7 gene expression.

#### Inactive Ribozyme Has *In Vivo* Antisense Activity

The effects of long-term expression of the E6/E7 genes with or without antisense containing ribozyme were investigated using transfected HKc as above, but maintained for up to 8 weeks in culture with standard cell culture medium after an initial G418 drug selection of four days. By the end of 8-week incubation, HKc transfected with a non-immortalizing gene (bacterial  $\beta$ -galactosidase) had senesced and detached. Therefore, mostly immortal cells are present in the cell cultures.

As shown in FIG. 7, cells transfected with the E6/E7 control (pCR16HH) were efficiently immortalized. Cells transfected with the active ribozyme construct (pCR16E6/E7RZ) showed little survival (about 10% relative to the control), as expected because of the limited E6/E7 gene expression. Cells transfected with the inactive ribozyme construct (pCR16E6/E7RZi) also showed decreased survival compared to the control, indicative that the antisense oligonucleotide portion of the construct, even in the absence of ribozyme activity, significantly inhibited E6/E7 gene expression. The lack of E6/E7 gene expression was confirmed using the RT-PCR assay which did not detect full length (492 bp) E6/E7 transcripts in either the cells transfected with active or inactive ribozyme constructs (pCR16E6/E7RZ or pCR16E6/E7RZi). These results also show the inhibitory activity *in vivo* of the antisense E6 oligonucleotide moiety of the pCR16E6/E7RZi construct.

To determine whether antisense oligodeoxynucleotides (ODN) complementary to HPV E6 could direct RNase-mediated degradation of the E6 mRNA, a  $^{32}$ P-labeled synthetic RNA target from HPV-16 (nucleotides 413-446 of SEQ ID NO: 1) was produced by *in vitro* transcription and incubated with 2 units *E. coli* RNase H (Life Technologies) and 10 nmoles of ODN, which were synthesized in an Expedite 8900 DNA synthesizer using phosphoramidite chemistry (Perseptive Biosystems): Anti-E6 (5'-TTCTTCAGGACACAGT-3'; SEQ ID NO: 18, complement of SEQ ID NO: 3), M4 (5'-TTCTTCAGAGAACAGT-3'; SEQ ID NO: 7) or M7 (5'-TTCTTACTAGAACAGT-3'; SEQ ID NO: 19) at 37°C for up to 30 min in RNase H buffer (20 mM Tris-HCl, pH 8.0, 100 mM KCl, 10 mM MgCl<sub>2</sub>, 0.1 mM EDTA, 0.1 M dithiothreitol). Anti-E6 is complementary to nucleotides 430-445 of HPV-16 E6 mRNA. The oligomer 5'-GGCTTTTGACAGTTA-3' (SEQ ID NO: 20) is complementary to the HPV-16 sequence from nt 415-429 (SEQ ID NO: 2). The oligomer M4 and M7 differ from Anti-E6 by 4 and 7 bases, respectively (Table 1). Additional experiments used  $^{32}$ P-labeled full-length HPV-16 E6/E7 genes (nt 97 to 868) under identical reaction conditions. Reactions were stopped with gel loading buffer (80% formamide, 0.01% bromphenol blue, 0.01% xylene cyanol) and heated at 65°C

for 5 min, then analyzed by electrophoresis in 7 M urea 6% polyacrylamide gels. Gels were dried and exposed to Kodak BioMax BM radiographic film (Eastman Kodak, Rochester, NY).

**Table 1. Synthetic ODN used. Mismatched bases are underlined.**

5	Name	Sequence
	Anti-E6	5'-TTC TTC AGG ACA CAG T-3'
	M4	5'-TTC TTC AGA <u>GAA</u> CAG T-3'
	M7	5'-TTC TTA <u>CTA GAA</u> CAG T-3'
10		

Anti-E6 and M4 ODN were able to direct RNase H-mediated degradation of the target mRNA as shown by specific degradation of the transcript visualized by urea polyacrylamide gel electrophoresis, while the M7 mutant failed to direct RNase activity. Anti-E6 directed RNase H activity over the full length HPV-16 E6/E7 transcript; however, neither M4 or M7 affected this target. Thus, the Anti-E6 antisense ODN effectively directs RNase H activity over the entire HPV-16 E6/E7 mRNA.

The *in vivo* efficacy of single-stranded ODN can be seriously compromised by cellular exo- and endonucleases. To estimate the *in vivo* survival rate of HPV-16 antisense ODN, a 5'-fluorescein labeled Anti-E6 ODN (Genosys Biotechnologies, Inc.) was purified by high performance liquid chromatography (HPLC) and applied to HPV-16 cultured immortal (Hkc 16E6/E7-II) and cervical tumor cell lines (CasKi; ATCC CRL-1550 and SiHa; ATCC HTB-35) and to normal human keratinocytes (Hkc) from neonatal foreskins. HKc were cultured in keratinocyte-SFM (Life Technologies). CasKi cells were cultured in DMEM (Life Technologies) enriched with 5% fetal bovine serum (FBS, Gemini Bio-Products). Cells were cultured in 8-well slide chambers (Nunc, Inc., Naperville, IL) until 70% confluent in the appropriate media. Fresh medium containing 10 mM 5-fluorescein labeled ODN was added and cells were incubated for 2 hours at 37°C. Cultures were washed twice with phosphate buffered saline (PBS) and fresh medium was added. The slides were further incubated for various times (0-16 hours) at 37°C. ODN survival was estimated by the number of fluorescent cells detected with a Leitz Ortholux II fluorescence microscope using a FITC filter. The results are summarized in Table 2.

**Table 2. Anti-E6 survival in different cell types. N, nuclear localization; C, cytoplasmic localization; ND, not detected; o/n, overnight.**

30	Cell type	0 h	1h	2h	o/n
	HKc	++++ N	++++ N	++++ N	++++ N
	HKc16E6/E7II	++++ N	+++ NC	++ NC	+ NC
35	SiHa	++++ N	++ NC	++ NC	+ NC
	CasKi	++++ N	+ N	ND	ND

The fluorescent label persisted for about 1 hour in tumor cell lines and for more than 2 hours in HPV-16 immortalized cell lines. No fluorescence remained after 12 hours. In contrast, normal keratinocytes retained the fluorescent label for 16 hours. These results suggest that the stability of the Anti-E6 ODN correlated with the abundance of the target (HPV-16 E6). Thus, it appears that the efficiency of antisense inhibition is dependent on the steady state level of the ODN.

Modification of the chemical structure of therapeutic ODNs has been shown to effectively increase their half-life *in vivo* and to retain their capacity to direct RNase H activity on the target RNA (Yamaguchi et al., *Leukemia* 11:497-503, 1997; Temsamani et al., *J. Biol. Chem.* 266:468-472, 1991; Boiziau et al., *Biochimie* 73:1403-1408, 1991; Akhtar et al., *Life Sci.* 49:1793-1801, 1991; Agrawal et al., *Proc. Natl. Acad. Sci. U.S.A.*, 87:1401-1405, 1990). CasKi and QGU cervical tumor cells were cultured in 96-well plates and starved for 48 hours before treatment with 50  $\mu$ M anti-E6 or M7 phosphorothioated ODNs (S-ODNs) for 72 hours. QGU cells (Shirasawa et al., *J. Gen. Virol.* 68:583-591, 1987) were cultured in F12/DMEM medium (Life Technologies). Anti-E6, M4 and M7 S-ODNs were obtained from Genosys Biotechnologies, purified by HPLC and applied to cells (50  $\mu$ M) for 72 hours. ODN attrition was compensated by adding fresh ODN containing medium every 24 hours. Cell growth was estimated by the colorimetric quantification of Br-dUTP incorporation using the Cell Proliferation Kit III (Boehringer Mannheim, Indianapolis, IN). Growth inhibition of CasKi cells was observed only by the wild-type Anti-E6 ODN (Fig. 8). Untreated serum-deprived CasKi cells were used as a negative control. Similar results were obtained for QGU cells.

The antisense oligodeoxynucleotides (SEQ ID NO:6, SEQ ID NO:7) corresponding to the antisense sequences contained in the ribozymes (SEQ ID NO:4, SEQ ID NO:5) are synthesized as normal phosphodiester bond-linked oligonucleotides and phosphorothioate oligonucleotides to inhibit E6/E7 gene expression in human cells infected with HPV. The phosphorothioate antisense oligonucleotides (PS-oligonucleotides) are synthesized using standard methods (Agrawal et al., 1997, *Proc. Natl. Acad. Sci. USA* 94:2620-2625; Agrawal, 1996, *TIBTECH* 14:376-387). That is, the oligonucleotides are synthesized in which one of the non-bridging oxygens of the internucleotide phosphodiester linkages is replaced with sulfur; the synthesis is done using methods that produce a diastereomeric mixture of Rp and Sp PS-oligonucleotides. HKC that have been shown to contain HPV sequences or suspected of containing HPV sequences due to their source of origin (e.g., cervical cancer cells) are then transfected with about 25 nM to about 500 nM concentration, preferably about 200 nM concentrations, of oligonucleotides and PS-oligonucleotides, using lipofection as described above or any well known transfection methodology. Cells of the same origin as those transfected are used as a source of purified RNA, essentially as described above. Transfected cells are allowed to grow for 2-10 days in culture and then RNA is similarly isolated from them. E6 transcripts are assayed in the untransfected control cells and the antisense-oligonucleotide and antisense PS-oligonucleotide transfected cells using the RT-PCR assay substantially as described above. The amount of E6 transcript detected in transfected and untransfected cells of the same origin are compared and quantified.

In non-transfected control cells, E6 transcripts are present in most cells known to harbor HPV sequences and in most cervical cancer cells isolated from invasive cancers. In the matched cells for each sample transfected

with the antisense oligonucleotides and antisense PS-oligonucleotides, there is measurably less E6 transcript detected compared to the matched control. The inhibition of E6 gene expression ranges from about 10% to about 95%, depending on the combination of the cells used and the antisense oligonucleotide or PS-oligonucleotide. Some cells show about 70% to about 80% inhibition, whereas others show about 40% to about 50% inhibition, and still others show about 10% to about 25% inhibition of E6 gene expression compared to the matched untransfected control cells. In all cells, a positive control (e.g.,  $\beta$ -actin gene expression (detected using the RT-PCR assay and primers of SEQ ID NO:14 and SEQ ID NO:15, as described above) varies by less than about 1% to about 5% from the HPV-antisense transfected cells and their matched untransfected control cells.

Similarly, antisense oligonucleotide and PS-oligonucleotide having sequences of SEQ ID NO:17 (TTCTTCAGAGAACAGTGGCTTTTGACAGTTA; corresponding to the antisense of the RNA sequence SEQ ID NO:16: UUCUUCAGAGAACAGUGGCUUUUGACAGUUA), representing the antisense DNA to nt 415 to 445 of HPV-16 are also synthesized using standard methods. Similarly the longer antisense oligonucleotide and PS-oligonucleotide are used in transfections of HKc, substantially as described above. The results of RT-PCR assays to determine the relative amounts of E6 transcripts in antisense oligonucleotide and PS-oligonucleotide transfected cells, compared to untransfected matched control cells, show somewhat less inhibition of E6 transcripts in the transfected cells compared to the smaller antisense oligonucleotide and PS-oligonucleotides described above. That is, about 0% to about 50% inhibition of E6 transcripts is detected in the tested cells. This lower level of inhibition may reflect less efficient transfection with the longer antisense oligonucleotide and PS-oligonucleotide.

When proliferation of the transfected cells (with any of the above-described antisense oligonucleotides and PS-oligonucleotides) and the matched untransfected control cells are compared using cell culture times of 1 to 5 days (starting at day 0, with about  $10^6$  cells/well of a standard 6-well culture plate and standard tissue culture conditions), the untransfected control cells generally show considerably more cell growth during the growth period than the matched transfected cells. The degree of proliferation inhibition is best seen at days 1 to 3, with some transfected cells showing approximately the same rate of cell proliferation as the matched control cells by day 5 after transfection. Some transfected cells show no inhibition of cell growth at any point in the testing period. Others show about 5% to about 40% inhibition of growth, depending on the cell line and the antisense oligonucleotide used. Retransfection at day 4, with those cells that initially show cell proliferation, reinstitutes the inhibition caused by the antisense oligonucleotides.

Similar inhibition results are obtained with antisense oligonucleotide sequences (SEQ ID NO:6, SEQ ID NO:7 and SEQ ID NO:17) synthesized as mixed backbone oligonucleotides, having 2'-O-methylnucleoside phosphodiester bonds in place of phosphorothioate bonds in some positions, and oligonucleotides synthesized with ends of 2'-O-methylnucleosides, both types synthesized using known methods (Agrawal et al., 1997, *Proc. Natl. Acad. Sci. USA* 94:2620-2625). The mixed backbone antisense oligonucleotides, synthesized as racemic mixtures, are preferable to completely PS-oligonucleotides because of reduced toxicity to cells exhibited by the mixed backbone oligonucleotides.

Similar tests are performed with the antisense oligonucleotide sequences (SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:17) synthesized as analogs using standard methods. The analogs synthesized are methylphosphonates

(Miller et al., 1993, in *Antisense Research and Applications*, pp.189-203, Crooke & Lebleu, eds., CRC Press; Sarin et al., 1988, *Proc. Natl. Acad. Sci. USA* 85:7448-7451); phosphoramidites (Dagle et al., 1991, *Nucl. Acids Res.* 19(8):1805-1810; Froehler et al., 1988, *Nucl. Acids Res.* 16(11):4831-4839; Tanaka et al., 1987, *Nucl. Acids Res.* 15(15):6209-6224); phosphorodithioates (Marshall et al., 1992, *Proc. Natl. Acad. Sci. USA* 89:6265-6269), and N3'→P5'-phosphoramidites (Gryznov et al., *Nucl. Acids Res.* 24:1508-1514; Escude et al., 1996, *Proc. Natl. Acad. Sci. USA* 93(9):4365-4369; Chen et al., 1995, *Nucl. Acids Res.* 23(14):2661-2668). Analogs of the antisense oligoribonucleotides (SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:16) are synthesized using known methods to produce oligoribonucleotide phosphorothioates (Agrawal et al., 1992, *Ann. New York Acad. Sci.* 660:2-10) and their 2'-O-alkyl analogs (Metelev et al., 1994, *Bioorg. Med. Chem. Lett.* 4:2929-2934; McKay et al., 1996, *Nucl. Acids Res.* 24:411-417; Monia et al., *J. Biol. Chem.* 268:14514-14522) and 2'-O-methylribonucleotide methylphosphonates (Kean et al., 1995, *Biochemistry* 34:14617-14620). All of the above-cited methods are known in the art and can readily be practiced by those skilled in the art; however, details regarding synthesis methods contained in the references cited herein are hereby incorporated by reference.

#### Antitumor Activity of Antisense Therapeutics in a Mouse Model

To determine whether antisense HPV-16 E6 ODN could inhibit the growth of cervical tumors *in vivo*, female C57 nude mice (3-4 weeks old) were injected subcutaneously with  $5 \times 10^6$  CasKi cells where they produce solid tumors or ascites. When tumors were palpable, Alzet osmotic pumps model 1002 (Alza Corporation, Mountain View, CA) filled with 500  $\mu$ g of Anti-E6 (SEQ ID NO: 18) or M7 (SEQ ID NO: 19) S-ODN in sterile water were implanted near the tumors for 14 days. Control animals received the tumor cells, but were treated with vehicle only. Anti-E6 inhibited the growth of these tumors, while M7 was ineffective. In M7 (SEQ ID NO: 19)-treated mice, tumor growth was identical to that observed with control animals. Thus, HPV-16 antisense oligonucleotides effectively inhibit cervical tumor cell growth in an *in vivo* animal model.

Other antisense oligonucleotides, antisense PS-oligonucleotides, mixed backbone antisense oligonucleotides, oligonucleotides having ends of 2'-O-methylnucleosides, and the oligonucleotide analogs or oligoribonucleotide analogs can also be tested in this mouse model. Typically, HPV-containing cervical tumor cells are injected into nude mice (e.g., subcutaneous at  $10^6$  to  $10^8$  cells/mouse). Preferred tumor cells are those isolated from spontaneous cervical carcinomas that are HPV positive (16 cell lines of this type are currently available for use). The mice are injected with 100  $\mu$ g to 1,000  $\mu$ g per day of the antisense therapeutic (i.p., s.c. or i.v.) before, simultaneously with or after injection of the human tumor cells (using protocols substantially as described in Skorski et al., *Proc. Natl. Acad. Sci. USA*, 1997, 94(8):3966-3971). The mice are monitored for survival times and, using standard assay techniques, for production of solid tumors and/or ascites at daily to weekly intervals from 1 day to 5 months post-injection of the tumor cells.

Injection of the antisense therapeutics one day before or simultaneous with injection of the tumor cells shows that some of the antisense therapeutics have protective effects and prevent tumor development that otherwise occurs in the untreated controls injected with tumor cells. That is, treatment of mice with some of the antisense therapeutics entirely inhibits tumor cell growth or significantly slows growth of tumors or ascites compared to control

mice that receive the same number of injected tumor cells but no antisense treatment. For those antisense therapeutics that show antitumor activity in preliminary tests, daily injections of 100, 300 and 900  $\mu\text{g}$  are subsequently tested to determine optimum antitumor activity and toxicity levels resulting from daily injections.

Injection of the antisense therapeutics at 1 day to 4 weeks after injection of the tumor cells shows that some of the antisense therapeutics are capable of limiting or eliminating tumors that otherwise grow in the untreated controls. Generally, antisense treatment is more effective, both in terms of long-term survival and inhibition of tumor cell proliferation, when treatment begins early after injection of tumor cells (i.e., within 1 to 2 weeks) and when mice receive repeated injections of the antisense therapeutic (e.g., daily or weekly dosages) during the test period. Also, the analogs (e.g., mixed backbone oligonucleotides and methylphosphonates, phosphoramidites, phosphorodithioates, or N3'→P5'-phosphoramidites) are generally more effective at lower concentrations (e.g., 100  $\mu\text{g}$  to 500  $\mu\text{g/day}$ ) than the corresponding unmodified antisense oligonucleotides.

In other animal models, mice showing spontaneous growth of cervical cancers are treated by s.c. injection or painting of the cervical tumor with the antisense therapeutics described herein. Some tumors treated with some of the antisense therapeutics show a decrease in tumor growth or remission of the tumor. In general, the antisense therapeutics that are mixed backbone oligonucleotides or other analogs (e.g., methylphosphonates, phosphoramidites, phosphorodithioates, or N3'→P5'-phosphoramidites) are generally more effective at lower concentrations (e.g., 25 nM to 100 nM/day) than the corresponding unmodified antisense oligonucleotides.

Similarly, analogs of antisense oligoribonucleotides of SEQ ID NO:4, SEQ ID NO:5, and SEQ ID NO:16, in which the analogs are oligoribonucleotide phosphorothioates, 2'-O-alkyl oligoribonucleotide phosphorothioates or 2'-O-methylribonucleotide methylphosphonates, are tested to antitumor activity in the assays described above. Some of the antisense oligoribonucleotide analogs also show significant antitumor activity *in vitro* and *in vivo*.

HPV-16 antisense molecules which inhibit cervical tumor cell growth in the mouse model discussed above are used for inhibition of human cervical tumor growth. In a preferred embodiment, for treatment of human cervical tumors, the cervix is injected daily with an antisense HPV-16 molecule such as Anti-E6 (SEQ ID NO: 18) S-ODN in a pharmaceutically acceptable carrier such as sterile PBS, in an amount between about 10 mg and 10,000 mg, more preferably between about 100 mg and 1,000 mg. Alternatively, the Anti-E6 (SEQ ID NO: 18) S-ODN is painted onto the cervix. The Anti-E6 S-ODN may also be applied to a ring of biocompatible material such as polypropylene or polyethylene, which is then placed around the cervix. These administration protocols all result in uptake of the ODN by the cervical tumor cells, inhibition of HPV-16, and inhibition of tumor cell growth.

While particular embodiments of the invention have been described in detail, it will be apparent to those skilled in the art that these embodiments are exemplary rather than limiting, and the true scope of the invention is defined by the claims that follow.



**WHAT IS CLAIMED IS:**

1.       Analog of antisense oligonucleotides comprising oligonucleotide sequences complementary to SEQ ID NO:2 or SEQ ID NO:3, wherein the analogs are phosphorothioate antisense oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond, mixed backbone antisense oligonucleotides in which  
5       at least one phosphodiester bond is replaced with a phosphorothioate bond and at least one phosphodiester bond is replaced with a 2'-*O*-methylnucleoside phosphodiester bond, end-modified analogs in which at least one end has a 2'-*O*-methylnucleotide moiety, methylphosphonates, phosphoramidites, phosphorodithioates, or N3'→P5'-phosphoramidites.
2.       Therapeutic compositions comprising any one of the analogs of antisense oligonucleotides of Claim 1, or mixtures thereof, in a pharmaceutically acceptable carrier.
- 10       3.       Therapeutic compositions of Claim 2, further comprising a ribozyme containing sequences complementary to SEQ ID NO:2 or SEQ ID NO:3.
4.       Analog of antisense oligonucleotides comprising sequences of SEQ ID NO:6, SEQ ID NO:7 or SEQ ID NO:17, wherein the analogs are phosphorothioate oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond, mixed backbone oligonucleotides in which at least one phosphodiester bond  
15       is replaced with a phosphorothioate bond and at least one phosphodiester bond is replaced with a 2'-*O*-methylnucleoside phosphodiester bond, end-modified oligonucleotides in which at least one end has a 2'-*O*-methylnucleotide moiety, methylphosphonates, phosphoramidites, phosphorodithioates, or N3'→P5'-phosphoramidites.
5.       Therapeutic compositions comprising any one of the analogs of antisense oligonucleotides of Claim 4, or mixtures thereof, in a pharmaceutically acceptable carrier.
- 20       6.       Analog of antisense oligoribonucleotides of SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:16, wherein the analogs are oligoribonucleotide phosphorothioates, 2'-*O*-alkyl oligoribonucleotide phosphorothioates or 2'-*O*-methylribonucleotide methylphosphonates.
7.       Therapeutic compositions comprising any one of the antisense oligoribonucleotide analogs of Claim 6, or mixtures thereof, in a pharmaceutically acceptable carrier.
- 25       8.       An antisense therapeutic composition comprising an oligonucleotide sequence of SEQ ID NO: 9.
9.       A method of preventing transformation of a living cell by HPV comprising the steps of:  
          providing an antisense therapeutic composition according to Claim 8;  
          providing a living cell capable of being transformed by HPV;  
          transfecting said living cell with the antisense therapeutic composition; and  
30       maintaining said living cell alive for sufficient time to inhibit expression of HPV gene E6.
10.       The method of Claim 9, wherein the living cell is a human keratinocyte.
11.       The method of Claim 9, wherein the living cell is a human cervical cell.
12.       A method of preventing transformation of a living cell by HPV comprising the steps of:  
          providing one or more antisense oligonucleotides having a nucleotide sequence(s) complementary  
35       to SEQ ID NO:2 and/or SEQ ID NO:3;  
          providing a living cell capable of being transformed by HPV;

transfecting said one or more antisense oligonucleotides into the living cell; and  
maintaining viability of the living cell for sufficient time to inhibit expression of HPV gene E6.

13. The method of Claim 12, wherein the living cell is a human keratinocyte.

14. The method of Claim 12, wherein the living cell is a human cervical cell.

5 15. A method of inhibiting expression of HPV gene E6 in a living cell comprising the steps of:  
providing one or more antisense oligonucleotides having a nucleotide sequence(s) complementary  
to SEQ ID NO: 2 and/or SEQ ID NO: 3;

providing a biological sample comprising living cells capable of being infected with HPV;

transfecting said one or more antisense oligonucleotides into said living cells; and

10 maintaining the viability of said living cells for sufficient time to allow inhibition of HPV E6 gene  
expression to occur.

16. The method of Claim 15, further comprising repeating the transfecting and maintaining steps.

17. The method of Claim 15, wherein the living cells are human keratinocytes.

18. The method of Claim 15, wherein the living cells are human cervical cells.

15 19. The method of Claim 15, wherein the step of providing antisense oligonucleotides comprises  
administering said one or more antisense oligonucleotides to a living organism by subcutaneous, intraperitoneal or  
intravenous injection, or by painting said antisense oligonucleotides onto said biological sample *in situ*.

20. A method of inhibiting expression of HPV gene E6 in a living cell comprising the steps of:  
providing one or more antisense oligonucleotide analogs having sequences of SEQ ID NO:6, SEQ  
20 ID NO:7, or SEQ ID NO:17, wherein the one or more antisense oligonucleotide analogs are phosphorothioate  
oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond, mixed  
backbone oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate  
bond and at least one phosphodiester bond is replaced with a 2'-O-methylnucleoside phosphodiester bond,  
in which at least one end has a 2'-O-methylnucleotide moiety, methylphosphonates, phosphoramidites,  
25 phosphorodithioates, or N3'→P5'-phosphoramidites;

providing a biological sample comprising living cells capable of being infected with HPV;

transfecting said one or more antisense oligonucleotide analogs into said living cells; and

maintaining the viability of said living cells for sufficient time to allow inhibition of HPV E6 gene  
expression to occur.

30 21. A method for inhibiting expression of HPV gene E6 in a living cell comprising the steps of:  
providing one or more antisense oligoribonucleotide analogs of oligoribonucleotides of SEQ ID NO:4,  
SEQ ID NO:5 or SEQ ID NO:16, wherein the one or more analogs are oligoribonucleotide phosphorothioates,  
2'-O-alkyl oligoribonucleotide phosphorothioates or 2'-O-methylribonucleotide methylphosphonates;

providing a biological sample comprising living cells capable of being infected with HPV;

35 transfecting said one or more antisense oligoribonucleotide analogs into said living cells; and

maintaining the viability of said living cells for sufficient time to allow inhibition of HPV E6 gene expression to occur.

22. A method for inhibiting the growth of a cervical tumor, comprising the step of contacting said tumor with one or more analogs of antisense oligonucleotides comprising oligonucleotide sequences complementary to SEQ ID NO:2 or SEQ ID NO:3, wherein the one or more analogs are phosphorothioate antisense oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond, mixed backbone antisense oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond and at least one phosphodiester bond is replaced with a 2'-*O*-methylnucleoside phosphodiester bond, end-modified analogs in which at least one end has a 2'-*O*-methylnucleotide moiety, methylphosphonates, phosphoramidites, phosphorodithioates, or N3'→P5'-phosphoramidites.

23. Use of one or more analogs of antisense oligonucleotides comprising oligonucleotide sequences complementary to SEQ ID NO:2 or SEQ ID NO:3, wherein the one or more analogs are phosphorothioate antisense oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond, mixed backbone antisense oligonucleotides in which at least one phosphodiester bond is replaced with a phosphorothioate bond and at least one phosphodiester bond is replaced with a 2'-*O*-methylnucleoside phosphodiester bond, end-modified analogs in which at least one end has a 2'-*O*-methylnucleotide moiety, methylphosphonates, phosphoramidites, phosphorodithioates, or N3'→P5'-phosphoramidites, for treatment of a cervical tumor.

1/8




FIG. 1

2/8

# HPV-16 E6 mRNA




FIG. 2

3/8




FIG. 3

4/8




FIG. 4

5/8




FIG. 5



6/8




FIG. 6

7/8




FIG. 7

8/8




FIG. 8

## SEQUENCE LISTING

5           < 110 > Government of the United States of America,  
             As Represented by the Secretary, Department  
             of Health and Human Services  
             DiPaolo, Joseph  
             Alvarez-Salas, Luis

10          < 120 > HUMAN PAPILLOMA VIRUS INHIBITION BY  
             ANTISENSE OLIGONUCLEOTIDES

15          < 130 > NIH138.001QPC

            < 150 > US 08/929,140  
             < 151 > 1997-09-05

20          < 160 > 20

            < 170 > FastSEQ for Windows Version 3.0

25          < 210 > 1  
             < 211 > 7904  
             < 212 > DNA  
             < 213 > Human Papilloma Virus 16

            < 400 > 1

	actacaataa ttcatgtata aaactaaggc cgtaaccgaa atcgggtgaa ccgaaaccgg	60
30	ttagtataaa agcagacatt ttatgcacca aaagagaact gcaatgttcc aggaccaca	120
	ggagcgacc agaaagttac cacagttagt cacagagctg caaacaacta tacatgat	180
	aatattagaa tgtgtgtact gcaagcaaca gttactgcga cgtgaggtat atgactttgc	240
	ttttcgggat ttatgcatag tatatagaga tgggaatcca tatgctgtat gtgataaatg	300
	tttaaagttt tattctaaaa ttagtgagta tagacattat tgttatagtt tgtatggaac	360
35	aacattagaa cagcaataca acaaaccgtt gtgtgatttg ttaattaggt gtattaactg	420
	tcaaaagcca ctgtgtcctg aagaaaagca aagacatctg gacaaaaagc aaagattcca	480

	taatataagg ggtcgggtgga ccggtcgaig tatgtcttgt tgcagatcat caagaacacg	540
	tagagaaacc cagctgtaat catgcatgga gatacaccta cattgcatga atatatgtta	600
	gatttgaac cagagacaac tgatctctac tgttatgagc aattaaatga cagctcagag	660
	gaggaggatg aaatagatgg tccagctgga caagcagaac cggacagagc ccattacaat	720
5	attgtaacct ttgttgcaa gtgtgactct acgcttcggt tgtgcgtaca aagcacacac	780
	gtagacattc gtactttgga agacctgta atgggcacac taggaattgt gtgccccatc	840
	tgttctcaga aaccataatc taccatggct gatcctgcag gtaccaatgg ggaagagggt	900
	acgggatgta atggatgggt ttatgtagag gctgtagtgg aaaaaaaac aggggatgct	960
	atatcagatg acgagaacga aaatgacagt gatacagggt aagatttgggt agattttata	1020
10	gtaaatgata atgattattt aacacaggga gaaacagaga cagcacatgc gttgtttact	1080
	gcacagggaag caaaacaaca tagagatgca gtacagggtc taaaacgaaa gtatttggtta	1140
	gtccacttag tgatattagt ggatgtgtag acaataatat tagtcctaga ttaaagcta	1200
	tatgtataga aaaacaaagt agagctgcaa aaaggagatt atttgaagc gaagacagcg	1260
	ggtagggcaa tactgaagtg gaaactcagc agatgttaca ggtagaaggg cgccatgaga	1320
15	ctgaaacacc atgtagtcat tatagtgggt gaagtggggg tggttgcagt cagtacagta	1380
	gtggaagtgg gggagagggt gttagtgaag gacacactat atgccaaca ccacttaca	1440
	atattttaa tgtactaaa actagtaatg caaaggcagc aatgttagca aaatttaag	1500
	agttatacgg ggtgagtttt tcagaattag taagaccatt taaaagtaat aaatcaacgt	1560
	gttgcgattg gtgtattgct gcatttgac ttacaccag tatagctgac agtataaaa	1620
20	cactattaca acaatattgt ttatatttac acattcaaag ttagcatgt tcatggggaa	1680
	tgggtgtgtt actattagta agatataaat gtggaaaaa tagagaaaca attgaaaaat	1740
	tgctgtctaa actattatgt gtgtctcaa tgtgtatgat gatagagcct ccaaattgc	1800
	gtagtacagc agcagcatta tattgttata aaacaggat atcaaatatt agtgaagtgt	1860
	atggagacac gccagaatgg atacaagac aaacagtatt acaacatagt ttaatgatt	1920
25	gtacattga attatcacag atgttacaat gggcctacga taatgacata gtagacgata	1980
	gtgaaattgc atataaatat gcacaattgg cagacactaa tagtaatgca agtgccttgc	2040
	taaaaagtaa ttcacaggga aaaattgtaa aggattgtgc aacaatgtgt agacattata	2100
	aacgagcaga aaaaaacaa atgagtatga gtcaatggat aaaatataga tgtgatagg	2160
	tagatgatgg aggtgattgg aagcaaattg ttatgtttt aaggatcaa ggtgtagagt	2220
30	ttatgtcatt ttaactgca ttaaaaagat ttttcaagg catacctaaa aaaaattgca	2280
	tattactata tgggtcagct aacacaggta aatcattatt tggatgagt ttaatgaaat	2340
	ttctgcaagg gtctgtaata tgtttgttaa attctaaaag ccatttttgg ttacaacat	2400
	tagcagatgc caaaatagggt atgttagatg atgctacagt gccctgttgg aactacatag	2460
	atgacaattt aagaaatgca ttggatggaa atttagtttc tatggatgta aagcatagac	2520
35	cattggtaca actaaaatgc cctccattat taattacatc taacattaat gctggtacag	2580
	attctagggt gccttattta cataatagat tgggtgtgtt tacatttcct aatgagtttc	2640

catttgacga aaacggaaat ccagtgtatg agcttaatga taagaactgg aaatcctttt 2700  
 tctcaaggac gtgggccaga ttaagtttgc acgaggacga ggacaaggaa aacgatggag 2760  
 actctttgcc aacgtttaaa tgtgtgtcag gacaaaatac taacacatta tgaaaatgat 2820  
 agtacagacc tacgtgacca tatagactat tggaaacaca tgcgcctaga atgtgtctatt 2880  
 5 tattacaagg ccagagaaat gggattttaa catattaacc accaagtggg gccaaacttg 2940  
 gctgtatcaa agaataaagc attacaagca attgaactgc aactaacgtt agaaacaata 3000  
 tataactcac aatatagtaa tgaaaagtgg acattacaag acgttagcct tgaagtgtat 3060  
 ttaactgcac caacaggatg tataaaaaaa catggatata cagtgggaagt gcagtttgat 3120  
 ggagacatat gcaatacaat gcattatata aactggacac atatatatat ttgtgaagaa 3180  
 10 gcatcagtaa ctgtggtaga gggtaagtt gactattatg gtttatatta tgttcatgaa 3240  
 ggaatagcaa catattttgt gcagttttaa gatgatgcag aaaaatatag taaaaataaa 3300  
 gtatgggaag ttcatgcggg tggtcaggta atattatgtc ctacatctgt gtttagcagc 3360  
 aacgaagtat cctctcctga aattattagg cagcacttgg ccaaccaccc cgccgcgacc 3420  
 cataccaaag cctgcgcctt gggcaccgaa gaaacacaga cgactatcca gcgaccaaga 3480  
 15 tcagagccag acaccggaaa cccctgccac accactaagt tgttcacag agactcagt 3540  
 gacagtgtc caatcctcac tgcatttaac agctcacaca aaggacggat taactgtaat 3600  
 agtaacacta caccatagat acatttataa ggtgatgcta atactttaa atgtttaaga 3660  
 tatagattta aaaagcattg tacattgtat actgcagtgt cgtctacatg gcattggaca 3720  
 ggacataatg taaaacataa aagtgaatt gttaactta catatgatag tgaatggcaa 3780  
 20 cgtgaccaat ttttgtcga agttaaata ccaaaaacta ttacagtgc tactggattt 3840  
 atgtctatat gacaaatctt gatactgcat ccacaacatt actggcgtgc ttttgcctt 3900  
 gctttgtgtg ctttgtgtg tctgcctatt aatacgtccg ctgctttgt ctgtgtctac 3960  
 atacacatca ttaataatat tggattact atgtggata acagcgcct ctgcgttag 4020  
 gtgttttatt gtatatatta tattgttta tataccatta ttttaatac atacacatgc 4080  
 25 acgtcttita attacataat gtatatgtac ataagttaatt tttacatat aattgttgta 4140  
 taccataact tactattttt tctttttat tttcatatat aattttttt tttgtttgtt 4200  
 tgtttgtttt ttaataaact gttattactt aacaatgcga cacaacgtt ctgcaaaacg 4260  
 caaaaaacgt gcatcggcta cccaacttta taaaacatgc aaacaggcag gtacatgtcc 4320  
 acctgacatt atacctaagg ttgaaggcaa aactattgct gaacaaatat tacaatatgg 4380  
 30 aagtatgggt gtattttttg gtgggttagg aattggaaca gggtcgggta caggcggagc 4440  
 cactgggtat attccattgg gaacaaggcc tcccacagct acagatacac ttgtcctgt 4500  
 aagacccctt ttaacagtag atcctgtggg ccttctgat cttctatag tttcttagt 4560  
 ggaagaaact agttttattg atgtgtgtgc accaacatct gtaccttcca ttccccaga 4620  
 tgtatcagga tttagtatta ctacttcaac tgataccaca cctgctatat tagatattaa 4680  
 35 taatactgtt actactgtta ctacacataa taatcccact ttcactgacc catctgtatt 4740  
 gcagcctcca acacctgcag aaactggagg gcattttaca ctttcatcat ccactattag 4800

	tacacataat tatgaagaaa ttcctatgga tacatttatt gtagcacaa accctaacac	4860
	agtaactagt agcacaccca taccagggtc tcgcccagtg gcacgcctag gattatatag	4920
	tcgcacaaca caacagggta aagttgtaga ccttgctttt gtaaccactc ccactaaact	4980
	tattacatat gataatcctg catatgaagg tatagatgtg gataatacat tatatttttc	5040
5	tagtaatgat aatagtatta atatagctcc agatcctgac tttttggata tagttgctti	5100
	acataggcca gcattaacct ctaggcgtac tggcattagg tacagtagaa ttggaataa	5160
	acaacacta cgtactcgtg gtggaaaatc tatagggtct aaggtagatt attattatga	5220
	tttaagtact attgatcctg cagaagaaat agaattacaa actataacac ctctacata	5280
	tactaccact tcacatgcag cctcacctac ttctattaat aatggattat atgatattta	5340
10	tgcagatgac ttattacag atacttctac aaccccggtg ccatctgtac cctctacatc	5400
	tttatcaggt tatattcctg caaatacaac aattcctttt ggtggtgcat acaatattcc	5460
	tttagtatca ggtcctgata taccatttaa tataactgac caagctcctt cattaattcc	5520
	tatagtcca gggctccac aatatacaat tattgctgat gcaggtgact ttattttaca	5580
	tcctagtatt tacatgttac gaaaacgacg taaacgttta ccatattttt tttcagatgt	5640
15	ctctttggct gcctagttag gccactgtct acttgcctcc tgtcccagta tctaagggtg	5700
	taagcacgga tgaatatgtt gcacgcacaa acatatatta tcatgcagga acatccagac	5760
	tacttgcagt tggacatccc tattttccta ttaaaaaacc taacaataac aaaatattag	5820
	ttcctaaagt atcaggatta caatacaggg tatttagaat acatttacct gacccaata	5880
	agtttgggtt tcttgacacc tcattttata atccagatac acagcggctg gtttggcct	5940
20	gtgtagggtt tgaggtaggt cgtggtcagc cattagggtt gggcattagt ggccatcctt	6000
	tattaaataa attggatgac acagaaaatg ctagtgtta tgcagcaaat gcagggtgtg	6060
	ataatagaga atgtatatct atggattaca aacaaacaca attgtgttta attggttgca	6120
	aaccacctat aggggaacac tggggcaaag gatcccatg taccaatgtt gcagtaaatc	6180
	caggtgattg tccaccatta gagttaataa acacagtat tcaggatggt gatatggttc	6240
25	atactggctt tgggtctatg gactttacta cattacaggc taacaaaagt gaagtccac	6300
	tggatatttg tacatctatt tgcaaatatc cagattatat taaaatggtg tcagaacat	6360
	atggcgacag cttatttttt tatttacgaa gggaacaaat gttgttaga catttattta	6420
	atagggtggt tactgttggg gaaaatgtac cagacgattt atacattaaa ggctctgggt	6480
	ctactgcaaa tttagccagt tcaaattatt ttctacacc tagtggttct atggttacct	6540
30	ctgatgccca aatattcaat aaaccttatt ggttacaacg agcacagggc cacaataatg	6600
	gcatttgttg gggtaaccaa ctatttgta ctgtgttga tactacacgc agtacaata	6660
	tgctattatg tgctgccata tctacttcag aaactacata taaaaatact aactttaagg	6720
	agtacctacg acatggggag gaatatgatt tacagtttat tttcaactg tgcaaaataa	6780
	ccttaactgc agacgttatg acatacatat attctatgaa ttccactatt ttggaggact	6840
35	ggaatttttg tctacaacct cccccaggag gcacactaga agatacttat aggtttgtaa	6900
	cccaggcaat tgcttgtaa aaacatacac ctccagcacc taaagaagat gatcccccta	6960

aaaaatacac tttttgggaa gtaaatttaa aggaaaagtt ttctgcagac ctagatcagt 7020  
 ttcctttagg acgcaaatTT ttactacaag caggattgaa ggccaaacca aaatttacat 7080  
 taggaaaaag aaaagctaca cccaccacct catctacctc tacaactgct aaacgcaaaa 7140  
 aacgtaagct gtaagtattg tatgtatgtt gaattagtgt tgtttgtgt gtatatgttt 7200  
 5 gtatgtgctt gtatgtgctt gtaaatatta agttgtatgt gtgtttgtat gtatggtata 7260  
 ataaacacgt gtgtatgtgt ttttaaatgc ttgtgtaact attgtgtcat gcaacataaa 7320  
 taaacttatt gtttcaacac ctactaattg tgtgtgggtt attcattgta tataaactat 7380  
 atttgcata tctgttttt gttttatata tactatattt tgtagcgcca ggcccatttt 7440  
 gtgacttcaa ccgaattcgg ttgcattgctt ttggcacaa aatgtgtttt tttaaatagt 7500  
 10 tctatgtcag caactatggt ttaaacttgt acgtttcctg ctggccatgc gtgccaatc 7560  
 cctgttttcc tgacctgcac tgcttgccaa ccattccatt gttttttaca ctgcactatg 7620  
 tgcaactact gaactactat gtacattgtg tcatataaaa taaatcacta tgcgccaacg 7680  
 ccttacatc cgctgttagg cacatatttt tggcttgttt taactaacct aattgcatat 7740  
 ttggcataag gtttaaacct ctaaggccaa ctaaagtca ccctagtcca tacatgaact 7800  
 15 gtgtaaaggt tagtcataca ttgttcattt gtaaaactgc acatgggtgt gtgcaaaccg 7860  
 attttgggtt acacatttac aagcaactta tataataata ctaa 7904

&lt;210&gt; 2

&lt;211&gt; 15

20

&lt;212&gt; RNA

&lt;213&gt; Human Papilloma Virus 16

&lt;400&gt; 2

uaacugucua aagcc

15

25

&lt;210&gt; 3

&lt;211&gt; 16

&lt;212&gt; RNA

&lt;213&gt; Human Papilloma Virus 16

30

&lt;400&gt; 3

acuguguccu gaagaa

16

&lt;210&gt; 4

35

&lt;211&gt; 15

&lt;212&gt; RNA



<213> Human Papilloma Virus 16

<400> 4

ggcuuuuaga aguuu

15

5

<210> 5

<211> 16

<212> RNA

<213> Human Papilloma Virus 16

10

<400> 5

uucuucagag aacagu

16

15

<210> 6

<211> 15

<212> DNA

<213> Human Papilloma Virus 16

<400> 6

20

ggcttttaga agtta

15

<210> 7

<211> 16

<212> DNA

25

<213> Human Papilloma Virus 16

<400> 7

ttcttcagag aacagt

16

30

<210> 8

<211> 61

<212> RNA

<213> Human Papilloma Virus 16

35

<400> 8

uucuucagag aacaguacca gagaaacaca cggacuucgg uccgugguau auuaccuggu

60

a 61

<210> 9  
<211> 61  
5 <212> RNA  
<213> Human Papilloma Virus 16

<400> 9  
uuuuuacagag aacaguacca gagcgucaca cggacuucgg uccgugguau auuaccuggu 60  
10 a 61

<210> 10  
<211> 60  
<212> RNA  
15 <213> Human Papilloma Virus 16

<400> 10  
ggcuuuuaga aguuuaccag agaaacacac ggacuucggg ccgugguaua uuaccuggua 60

20 <210> 11  
<211> 18  
<212> DNA  
<213> Human Papilloma Virus 16

25 <400> 11  
cagcaatata acaaaccg 18

<210> 12  
<211> 16  
30 <212> DNA  
<213> Human Papilloma Virus 16

<400> 12  
cacgtagaaa cccagc 16  
35 <210> 13

&lt;211&gt; 22

&lt;212&gt; DNA

&lt;213&gt; Human Papilloma Virus 16

5

&lt;400&gt; 13

tagattatgg tttctgagaa ca

22

&lt;210&gt; 14

&lt;211&gt; 31

10

&lt;212&gt; DNA

&lt;213&gt; Homo Sapien

&lt;400&gt; 14

tgacggggtc acccacactg tgcccatct a

31

15

&lt;210&gt; 15

&lt;211&gt; 31

&lt;212&gt; DNA

&lt;213&gt; Homo Sapien

20

&lt;400&gt; 15

ctaagaagca ttgcggtgg acgatggagg g

31

&lt;210&gt; 16

25

&lt;211&gt; 31

&lt;212&gt; RNA

&lt;213&gt; Human Papilloma Virus 16

&lt;400&gt; 16

30

uucuucagag aacaguggcu uuugacaguu a

31

&lt;210&gt; 17

&lt;211&gt; 31

&lt;212&gt; DNA

35

&lt;213&gt; Human Papilloma Virus 16

	<400> 17	
	ttcttcagag aacagtggct ttgacagtt a	31
	<210> 18	
5	<211> 16	
	<212> DNA	
	<213> Human Papilloma Virus 16	
	<400> 18	
10	ttcttcagga cacagt	16
	<210> 19	
	<211> 16	
	<212> DNA	
15	<213> Human Papilloma Virus 16	
	<400> 19	
	ttcttactag aacagt	16
	<210> 20	
20	<211> 15	
	<212> DNA	
	<213> Human Papilloma Virus 16	
	<400> 20	
25	ggcttttgac agtta	15

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/18320

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/11 C07H21/00 A61K31/70 //C07K14/025,C12N9/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07H A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 95 31552 A (US HEALTH ;NORTHERN ILLINOIS UNIVERSITY (US); HAMPEL ARNOLD (US);) 23 November 1995 see abstract see page 20, line 1 - line 5 see claims 1,2 seq. IDs 3,6,8,11 ---	1-23
Y	WO 93 20095 A (ISIS PHARMACEUTICALS INC ;CROOKE STANLEY T (US); MIRABELLI CHRISTO) 14 October 1993 see abstract see page 24, line 17 - line 23 see example 3 see claims 1-9 --- -/--	1-23

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

23 February 1999

Date of mailing of the international search report

02/03/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Gall, I

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/18320

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 13833 A (GENTA INC) 26 May 1995 see abstract see examples 1-27 see examples A-H -----	1-23
A	WO 96 39501 A (HOFFMANN LA ROCHE ;HYBRIDON INC.(US)) 12 December 1996 see abstract see table 18 see example 5 -----	1-23

# INTERNATIONAL SEARCH REPORT

...ernational application No.

PCT/US 98/ 18320

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 19, 22, 23 and (as far as methods in vivo are envisaged) claims 9-18,20,21 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/18320

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9531552 A	23-11-1995	US 5683902 A	04-11-1997
		AU 2588795 A	05-12-1995
		CA 2190102 A	23-11-1995
		EP 0759992 A	05-03-1997
		JP 10500307 T	13-01-1997
WO 9320095 A	14-10-1993	US 5457189 A	10-10-1995
		AU 671630 B	05-09-1996
		AU 3943893 A	08-11-1993
		CA 2132424 A	14-10-1993
		EP 0637316 A	08-02-1995
		FI 944523 A	29-09-1994
		HU 69934 A	28-09-1995
		JP 2737894 B	08-04-1998
		JP 9117291 A	06-05-1997
		JP 2609424 B	14-05-1997
		JP 7501709 T	23-02-1995
		NO 943198 A	29-08-1994
		US 5665580 A	09-09-1997
		US 5681944 A	28-10-1997
		US 5756282 A	26-05-1998
WO 9513833 A	26-05-1995	AU 678085 B	15-05-1997
		AU 1181995 A	06-06-1995
		AU 687492 B	26-02-1998
		AU 1291595 A	06-06-1995
		AU 689182 B	26-03-1998
		AU 1291695 A	06-06-1995
		CA 2176498 A	26-05-1995
		EP 0729474 A	04-09-1996
		EP 0735899 A	09-10-1996
		EP 0743859 A	27-11-1996
		JP 9507836 T	12-08-1997
		JP 9505306 T	27-05-1997
		JP 9506248 T	24-06-1997
		NZ 276956 A	27-04-1998
		NZ 277616 A	19-12-1997
		NZ 277617 A	26-01-1998
		WO 9514030 A	26-05-1995
		WO 9513834 A	26-05-1995
		US 5792615 A	11-08-1998
		AU 2584395 A	16-11-1995
		WO 9528942 A	02-11-1995
WO 9639501 A	12-12-1996	AU 6300296 A	24-12-1996
		CA 2226457 A	12-12-1996
		EP 0832214 A	01-04-1998